



# The relationship between mantle pH and the deep nitrogen cycle

Sami Mikhail<sup>a,b,\*</sup>, Peter H. Barry<sup>c</sup>, Dimitri A. Sverjensky<sup>d</sup>

<sup>a</sup> *The School of Earth and Environmental Sciences, The University of St. Andrews, St. Andrews, UK*

<sup>b</sup> *St Andrews Centre for Exoplanet Science, The University of St. Andrews, UK*

<sup>c</sup> *The Department of Earth Sciences, The University of Oxford, Oxford, UK*

<sup>d</sup> *The Department of Earth and Planetary Sciences, Johns Hopkins University, Baltimore, MD, USA*

Received 31 August 2016; accepted in revised form 6 April 2017; available online 18 April 2017

## Abstract

Nitrogen is distributed throughout all terrestrial geological reservoirs (i.e., the crust, mantle, and core), which are in a constant state of disequilibrium due to metabolic factors at Earth's surface, chemical weathering, diffusion, and deep N fluxes imposed by plate tectonics. However, the behavior of nitrogen during subduction is the subject of ongoing debate. There is a general consensus that during the crystallization of minerals from melts, monatomic nitrogen behaves like argon (highly incompatible) and ammonium behaves like potassium and rubidium (which are relatively less incompatible). Therefore, the behavior of nitrogen is fundamentally underpinned by its chemical speciation. In aqueous fluids, the controlling factor which determines if nitrogen is molecular ( $N_2$ ) or ammoniac (inclusive of both  $NH_4^+$  and  $NH_3^0$ ) is oxygen fugacity, whereas pH designates if ammoniac nitrogen is  $NH_4^+$  or  $NH_3^0$ . Therefore, to address the speciation of nitrogen at high pressures and temperatures, one must also consider pH at the respective pressure–temperature conditions. To accomplish this goal we have used the Deep Earth Water Model (DEW) to calculate the activities of aqueous nitrogen from 1–5 GPa and 600–1000 °C in equilibrium with a model eclogite-facies mineral assemblage of jadeite + kyanite + quartz/coesite (metasediment), jadeite + pyrope + talc + quartz/coesite (metamorphosed mafic rocks), and carbonaceous eclogite (metamorphosed mafic rocks + elemental carbon). We then compare these data with previously published data for the speciation of aqueous nitrogen across these respective P-T conditions in equilibrium with a model peridotite mineral assemblage (Mikhail and Sverjensky, 2014). In addition, we have carried out full aqueous speciation and solubility calculations for the more complex fluids in equilibrium with jadeite + pyrope + kyanite + diamond, and for fluids in equilibrium with forsterite + enstatite + pyrope + diamond.

Our results show that the pH of the fluid is controlled by mineralogy for a given pressure and temperature, and that pH can vary by several units in the pressure-temperature range of 1–5 GPa and 600–1000 °C. Our data show that increasing temperature stabilizes molecular nitrogen and increasing pressure stabilizes ammoniac nitrogen. Our model also predicts a stark difference for the dominance of ammoniac vs. molecular and ammonium vs. ammonia for aqueous nitrogen in equilibrium with eclogite-facies and peridotite mineralogies, and as a function of the total dissolved nitrogen in the aqueous fluid where lower N concentrations favor aqueous ammoniac nitrogen stabilization and higher N concentrations favor aqueous  $N_2$ .

Overall, we present thermodynamic evidence for nitrogen to be reconsidered as an extremely dynamic (chameleon) element whose speciation and therefore behavior is determined by a combination of temperature, pressure, oxygen fugacity, chemical activity, and pH. We show that altering the mineralogy in equilibrium with the fluid can lead to a pH shift of up to 4 units at 5 GPa and 1000 °C. Therefore, we conclude that pH imparts a strong control on nitrogen speciation, and thus N flux, and should be considered a significant factor in high temperature geochemical modeling in the future. Finally, our modelling

\* Corresponding author at: The School of Earth and Environmental Sciences, The University of St. Andrews, St. Andrews, UK.  
E-mail address: [sm342@st-andrews.ac.uk](mailto:sm342@st-andrews.ac.uk) (S. Mikhail).

demonstrates that pH plays an important role in controlling speciation, and thus mass transport, of Eh-pH sensitive elements at temperatures up to at least 1000 °C.

© 2017 Elsevier Ltd. All rights reserved.

*Keywords:* Mantle pH; Nitrogen geochemistry; Deep nitrogen cycle; Mantle volatiles; Subduction zones; Mantle redox

## 1. INTRODUCTION

Nitrogen is the dominant gas in Earth's atmosphere and is distributed in all terrestrial geological reservoirs, which are in a constant state of disequilibrium (see Busigny and Bebout, 2013; Bebout et al., 2013, 2016; Johnson and Goldblatt, 2015). The flux of nitrogen between the surface and interior is governed by volcanism (out-gassing) and subduction (in-gassing); this interplay ultimately controls atmospheric N<sub>2</sub> levels (see discussion in Barry and Hilton, 2016), which are intimately linked with biology (Stüeken et al., 2016; Zerkle and Mikhail, 2017). But to constrain the N flux and/or the partial pressure of atmospheric N over geologically-long (>Ga) timescales is difficult due to a lack of samples (e.g. Som et al., 2012, 2016; Marty et al., 2013). Consequently, in order to address these issues, workers must combine the evidence from sparse deep-time N datasets with experimental and theoretical (thermodynamic) models to estimate past nitrogen dynamics.

At present, there is no consensus for the direction (i.e., positive or negative) or magnitude of the global N flux out of the Earth through time (Busigny et al., 2011; Busigny et al., 2013; Dauphas and Marty, 1999; Fischer et al., 2002; Marty and Dauphas, 2003; Busigny et al., 2003; Elkins et al., 2006; Philippot et al., 2007; Yokochi et al., 2009; Mohapatra et al., 2009; Halama et al., 2010, 2014; Palot et al., 2012; Mikhail et al., 2014; Mikhail and Sverjensky, 2014; Barry and Hilton, 2016; Zerkle and Mikhail, 2017). As a result, there is no quantitative explanation for the discrepancy between calculated N-fluxes from different arc systems (see recent reviews by Busigny and Bebout, 2013; Bebout et al., 2013, 2016), and thus the nature of the deep nitrogen cycle remains a controversial topic (Zerkle and Mikhail, 2017).

To understand the pathways followed by N during subduction requires a first-order understanding of the partitioning behavior of nitrogen within specific minerals. For example, molecular nitrogen (N<sub>2</sub>) and ammonia (NH<sub>3</sub>) are both neutrally charged, and therefore highly incompatible in most mineral phases (Brooker et al., 2003). Conversely, ammonium (NH<sub>4</sub><sup>+</sup>) is positively charged and has an ionic radius between those of Rb<sup>+</sup> and K<sup>+</sup>, and thus should be compatible in K-bearing phases such as phengite, phlogopite, K-bearing clinopyroxene, K-hollandite, Phase-X ((K)Mg<sub>2</sub>Si<sub>2</sub>O<sub>7</sub>H) (Honma and Itihara, 1983; Haendel et al., 1986; Busigny et al., 2003; Yokochi et al., 2009; Palya et al., 2011; Bebout et al., 2016). Ammonium has also been shown to dissolve as a trace component in K-absent mafic minerals (Watenphul et al., 2010; Li et al., 2013). Therefore, the behavior of nitrogen is predictably governed by the speciation of nitrogen where neutrally charged compounds are – thermodynamically speaking – highly incom-

patible (Blundy and Wood, 2003), an assertion that has been demonstrated experimentally (Brooker et al., 2003).

For high temperature aqueous systems, the speciation of N has been constrained as a function of oxygen fugacity (Mikhail and Sverjensky, 2014; Li and Keppler, 2014), and both studies agree that most nitrogen in upper mantle fluids should be ammoniac. The difference between these datasets is the nature of the ammoniac nitrogen present. For example, at 5 GP, 1000 °C, and an oxygen fugacity of –1 log units relative to the Quartz-Fayalite-Magnetite buffer reaction (QFM) the speciation of nitrogen in equilibrium with olivine (Fo<sub>90</sub>) was determined to be ammonia in quenched run products (using FTIR; Li and Keppler, 2014), but the predicted speciation of nitrogen in equilibrium with forsterite + enstatite is ammonium (Mikhail and Sverjensky, 2014). We note that this could be the result of quenching, where ammonium (a reactive ion) re-equilibrates to a stable state during cooling (pre-analysis). But the controlling factor, which designates if ammoniac nitrogen is NH<sub>4</sub><sup>+</sup> and NH<sub>3</sub> is not oxygen fugacity, but rather pH – and more alkaline conditions favor NH<sub>3</sub>. This implies that the pH of the fluid in equilibrium with olivine (Fo<sub>90</sub>; Li and Keppler, 2014) may be more alkaline than is predicted for pH of the fluid in equilibrium with forsterite + enstatite (where pH can be expressed as proportional to  $\log[\alpha_{\text{Mg}^{2+}}/(\alpha_{\text{H}^+})^2]$ ; Mikhail and Sverjensky, 2014). Overall, this result implies pH is a significant variable at high temperatures.


Here we investigate the effect of pH on the speciation of nitrogen under upper mantle P-T-/O<sub>2</sub> conditions as a function of system stoichiometry using a hypothetical host-rock mineralogy. As is the case with /O<sub>2</sub>, the pH of the system is governed by pressure, temperature and composition (P-T-X). In the case of nitrogen geochemistry for our study, this can be considered as the equilibrium between a fluid and a solid. Therefore if all else is equal, the stoichiometry of the system can exert control on the pH and therefore the speciation and behavior of nitrogen. This is because different protonation reactions can buffer the pH of the fluid to different values (e.g., whether the protonation reactions are governed by the  $\log[\alpha_{\text{Mg}^{2+}}/(\alpha_{\text{H}^+})^2]$  or  $\log[\alpha_{\text{Na}^+}/\alpha_{\text{H}^+}]$ ). Herein we investigate the role of a hypothetical mineralogy (for stoichiometry) on fluid pH by presenting the outputs from a series of thermodynamic calculations, and we discuss the implications of our findings in light of the deep-Earth nitrogen cycle, and mantle geochemistry.

## 2. METHODS

The speciation of aqueous ions, metal complexes, neutral species scrapped can be predicted by applying the Helgeson-Kirkham-Flowers (HKF) equations of state

(Helgeson and Kirkham, 1974a,b, 1976; Helgeson et al., 1981; as revised in Tanger and Helgeson, 1988, Shock and Helgeson, 1988). Prior to recent pioneering work (Pan et al., 2013; Facq et al., 2014; Sverjensky et al., 2014a) there was a historic limitation of pressure <0.5 GPa, due to the fact that the dielectric constant of water was not known at higher pressures (Johnson et al., 1992; Shock et al., 1992). This precluded the application of the HKF equations of state to matters concerning aqueous fluids in deep Earth systems (e.g., lower crust, upper mantle, and subduction zones). However, the dielectric constant of water has recently been constrained  $\leq 6$  GPa (Sverjensky et al., 2014a), which enables an extension of the P-T range for the application of the HKF equations of state for aqueous species up to  $\leq 6$  GPa and  $\leq 1200$  °C (Pan et al., 2013; Facq et al., 2014; Sverjensky et al., 2014a). As a result, it is now feasible to determine the speciation of aqueous ions, metal complexes, neutral species in the presence of minerals across conditions akin to the pressure and temperature pathway followed by a subducted-slab from Earth's surface to a depth of approximately 150 km using the Deep Earth Water (DEW) model. As previously described (Mikhail and Sverjensky, 2014; Sverjensky et al., 2014b; Sverjensky and Huang, 2015), the DEW model enables calculation of equilibrium constants involving aqueous species of all kinds as a function of temperature and pressure, and the incorporation of these equilibrium constants into aqueous speciation, solubility, and chemical mass transfer codes.

We have predicted the speciation of nitrogen as a function of P-T- $f\text{O}_2$ -pH. Note, pH is a function of the activities of  $\text{Mg}^{2+}$  and  $\text{Na}^+$  in supercritical aqueous fluid in equilibrium with specific mineral assemblages (Sverjensky et al., 2014a). The pH values denoted by subducted oceanic crust in Fig. 1a–d were constrained by equilibrium with model eclogite-facies mineral assemblages (jadeite + kyanite +  $\text{SiO}_2$  representing metasediment, and jadeite + pyrope + talc +  $\text{SiO}_2$  representing metamorphosed mafic rocks) at pressures of 1 and 5 GPa and temperatures of 600–1000 °C. The equilibrium between fluid and jadeite, kyanite,  $\text{SiO}_2$  enables derivation of the following equations:




$$\text{for which } \log K_2 = \log \frac{a_{\text{Na}^+}^2}{a_{\text{H}^+}^2} \quad (2)$$

Assuming pure minerals and unit activities of the minerals and the water results in

$$\text{pH} = \frac{1}{2} \log K_2 - \log a_{\text{Na}^+} \quad (3)$$

For the mafic eclogite-facies mineral assemblages, equilibrium between the components of jadeite, garnet, talc and  $\text{SiO}_2$  enables writing



$$\text{for which } \text{pH} = \frac{1}{2} \log K_5 - \log a_{\text{Na}^+} \quad (5)$$

It should be noted that  $\text{SiO}_2$  is coesite at 5 GPa (600 & 1000 °C), but at 1 GPa  $\text{SiO}_2$  is stable as either  $\alpha$ -Qtz (600 °C) and  $\beta$ - $\text{SiO}_2$  (1000 °C). A range of values for the  $a_{\text{Na}^+}$  from 0.01 to 1.0 was used in Eqs. (3) and (5) resulting in a range of estimated pH values. For example, at 5 GPa and 600 °C, the calculated pH values from Eqs. (3) and (5) are 3.1–5.1 and 3.2–5.2, for metasedimentary and mafic eclogites respectively. Because these two ranges are indistinguishable on this scale, we only plot the values from Eq. (3) in Fig. 1a–d (subducted oceanic crust). The assumption of pure minerals introduces small uncertainties on the scale of the plots shown. For example, a decrease in the activity of jadeite in Eq. (3) from 1.0 to 0.5 would produce a decrease in the pH in Eq. (5) of 0.30, which is relatively small compared to the range of  $\text{Na}^+$  activities assumed. Very low activities of jadeite would produce a bigger effect (e.g., if the activity of jadeite were reduced from 1.0 to 0.10, the pH would be decreased by 1.0 unit). Furthermore, by applying values of  $\log K$  calculated as described above, together with the range of activities for the  $\text{Mg}^{2+}$  and  $\text{Na}^+$  corresponds to a range of pH values for the fluids. It should be emphasized that uncertainties caused by using the pure minerals and assuming that the activity of water is unity are rather small on a logarithmic scale such as in Fig. 1a–d compared to the range of pH associated with the range of activities for either  $\text{Mg}^{2+}$  or  $\text{Na}^+$ .

The boundaries between the various nitrogen species shown in Fig. 1a–d depend on the stoichiometry of the equilibria, the magnitudes of the relevant equilibrium constants, and the total dissolved nitrogen concentration (Mikhail and Sverjensky, 2014). The latter dependence is a consequence of equilibria between  $\text{N}_2$ , which has two moles of N, and reduced N-species, which have only one mole of N in each. As a consequence, the speciation of N in aqueous fluids at equilibrium is a function of the total dissolved nitrogen in the fluid. In our calculations, we consider a large range of possible nitrogen concentrations from 0.001 to 1 m N, corresponding to concentrations between 14 ppm by mass and 1.4 wt.%. This range was selected in order to simulate a wide range of likely fluid compositions that might be found in nature, but also to constrain the effect of N concentration (order of magnitude scales).

Although nitrogen concentrations in aqueous fluids from the upper mantle are poorly known (because aqueous fluids from the upper mantle are rarely sampled), the concentrations in minerals are better constrained. We cite a range of nitrogen concentrations measured in mantle rocks and minerals to demonstrate the possibility that aqueous fluids in the mantle might also exhibit a wide range of nitrogen concentrations. For example, typical measurements of volcanic xenoliths and basaltic samples are approximately 0.1–10 ppm by mass (whole rock from Marty, 1995, mineral separates from Fischer et al., 2005; basaltic glasses from Barry et al., 2012), and the phlogopites ranges from 7.6 to 25.7 ppm by mass (Yokochi et al., 2009) and lithospheric and sublithospheric diamonds contain between <1 and >10,000 ppm by mass N (Smart et al., 2011; Mikhail et al., 2014). In fact, some very rare mantle xenoliths and diamonds contain fluid-inclusions of pure  $\text{N}_2$  (Andersen et al., 1995; Smith et al., 2014). Despite the fact that nitro-




Fig. 1. Calculated  $\log f_{\text{O}_2}$ -pH diagrams for nitrogen speciation in supercritical aqueous fluids using the Deep Earth Water (DEW) model (see Section 2). The colored boxes show where predicted compositions of upper mantle (peridotite; green), mantle wedge (arc peridotite; blue) and subducted oceanic crust (eclogite; orange) are predicted to plot under these  $f_{\text{O}_2}$  - pH conditions. The boundaries between nitrogen-species represent a range of total dissolved nitrogen. For the purposes of examining nitrogen speciation under redox conditions appropriate to silicate mantles with a peridotitic bulk composition, we have expressed the  $\log f_{\text{O}_2}$  relative to the quartz-fayalite-magnetite mineral buffer (expressed as  $\Delta\text{QFM}$  in log units). The range of  $f_{\text{O}_2}$  values for eclogites shown above represents the global range, and is not intended to represent any single geographical locality. The fields for the oxidation state of peridotitic mantle domains and eclogitic mantle domains are described in the text. The pH values represent the equilibrium with jadeite + kyanite +  $\text{SiO}_2$  and a range of total dissolved Mg concentrations (see Section 2). (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

gen is a trace component in material from shallow depths, typically sampled by volcanoes (see Johnson and Goldblatt et al., 2015), there are processes (i.e. metasomatism and melting) that can generate localized nitrogen-enrichment and render nitrogen a major volatile element in the mantle. Therefore, the concentrations used here (ca. 14 ppm by mass to 1.4 wt.%) are not unrealistic, and enable us to investigate the speciation of nitrogen as a function of P, T,  $f_{\text{O}_2}$ , pH, and nitrogen concentration.

### 3. RESULTS

Superposed on Fig. 1a-d are fields representing the calculated (theoretical and empirical) oxygen fugacities of peridotitic (Wood and Virgo, 1989; Canil et al., 1990; Ionov and Wood, 1992; Woodland et al., 1992, 1996, 2006; Brandon and Draper, 1996; Frost and McCammon, 2008) and eclogitic rocks (Simakov, 2006; Stagno et al., 2015) as well as the predicted range of pH values for aqueous fluids in equilibrium with hypothetical mineral assemblages representing model peridotite (from Mikhail and Sverjensky, 2014) and eclogite-facies mineral assemblages (this study).

The oxygen fugacities for peridotites in arc-mantle wedges (QFM to QFM + 2) vs. the oxygen fugacity of peridotites from sub-cratonic lithospheric mantle (QFM to QFM - 3) are taken from Frost and McCammon (2008).

The available data suggest eclogitic rocks cover a similar  $f_{\text{O}_2}$  range to their peridotitic counterparts (Simakov, 2006; Stagno et al., 2015; Smart et al., in press). We plot a larger range for the  $f_{\text{O}_2}$  of eclogites in the figures (QFM - 4) to account for the observed occurrence of carbides, nitrides and native metals in samples from the obducted Tibetan ophiolites ( $\leq\text{QFM} - 4$ ; Dobrzynetskiy et al., 2009). In addition, the sediments subducted beneath the Cyclades Greek islands are relatively oxidized ( $\geq\text{QFM} - 1$ ; Ague and Nicolescu, 2014), having stable carbonate phases. The range of  $f_{\text{O}_2}$  values for eclogites shown in Figs. 1a-d represents the global range, and is not intended to represent any single geographical locality (i.e., data from Tibetan and Greek obducted rocks are provided to illustrate the large range of  $f_{\text{O}_2}$  values in measured subducted assemblages).

The pH values plotted here correspond to the calculated equilibrium between water and pure forsterite + enstatite (model peridotite – Mikhail and Sverjensky, 2014) and hypothetical eclogite-facies mineral assemblages; jadeite + kyanite +  $\text{SiO}_2$  (metasediment – this study), jadeite + pyrope + talc +  $\text{SiO}_2$  (metamorphosed mafic rocks – this study), and carbonaceous eclogite-facies mineral assemblages (metamorphosed mafic rocks + elemental carbon; this study). Importantly, the relative proportions of the pure minerals do not influence the fluid chemistry, because the fluid chemistry is set by the equilibrium constants (see Section 2). In our model, pH is expressed by the  $\alpha_{\text{H}^+}$  (fluid)

and  $\alpha_{\text{Na}^+}$  (eclogite – this study) and  $\alpha_{\text{Mg}^{2+}}$  (peridotite; Mikhail and Sverjensky, 2014) (Eqs. (3) and (5)). We have applied a range of activities for  $\alpha_{\text{Na}^+}$  and  $\alpha_{\text{Mg}^{2+}}$  from 0.1 to 1 (e.g. Eq. (2)). Our calculations show the speciation of aqueous ammoniac nitrogen in equilibrium with eclogite-facies metasediments and mafic rocks (herein collectively referred to as eclogite-facies mineral assemblage; this study) is predicted to differ from aqueous nitrogen in equilibrium with peridotites under most of the conditions investigated (Fig. 1a–d). Under most conditions, ammonium ( $\text{NH}_4^+$ ) dominates in peridotite, and ammonia ( $\text{NH}_3^0$ ) is dominant in the eclogite-facies mineral assemblages; this is due to the eclogite-facies mineral assemblages buffering the fluid to higher pH conditions (Fig. 1a–d).

For an oxygen fugacity of QFM -1 to -2, 600 °C, and 1 or 5 GPa (Fig 1a and c respectively), the dominant nitrogen species in the fluid are predicted to be  $\text{NH}_3^0$  or  $\text{NH}_4^+$ , depending on pressure and pH. At 1000 °C and 1 or 5 GPa (Fig 1b and d respectively), the dominant nitrogen species in the fluid is predicted to be  $\text{N}_2^0$ . This temperature range is relevant to modern arc systems. For example, Syracuse et al. (2010) calculated the thermal models for 204 slab temperatures in arc systems on the Pacific rim and found the range to be 301 to 987 °C, with a mean of  $789 \pm 76$  °C. Thus, our results strongly suggest that the nitrogen speciation in eclogitic fluids in ‘cold’ subduction zones differs from those in ‘hot’ subduction zones. Note, the exact thermal regimes for hot and cold subduction zones are difficult to constrain, because there are a number of dependent and independent variables to consider which control the thermal-depth status of a subducted slab (age of slab, slab-dip, slab T beneath arc, Moho T beneath arc, velocity of subduction) and the temperature also varies significantly with distance from slab surface (in both directions; Syracuse et al., 2010). Furthermore, because geothermal gradients during subduction are non-linear, there are multiple degrees of freedom to explore. To examine this further, we direct the reader to Syracuse et al. (2010). Herein, we refer to cold subduction zones as those with a thermal gradient (at the slab surface) of  $<10$  °C/km.

A significant difference between cold and hot subduction regimes are shown more distinctly in Figs. 2a–b and 3a–b. We calculated a full aqueous speciation model at 5.0 GPa between 600 and 1000 °C at QFM - 2 for aqueous nitrogen in fluids in equilibrium with a model eclogite consisting of jadeite + pyrope + kyanite + coesite, +diamond and a model peridotite consisting of forsterite + enstatite + pyrope + diamond. The latter is a more complex chemical system than has previously been considered (Mikhail and Sverjensky, 2014). Results are shown for high (0.1 m N  $\approx$  1.4 wt% N) and low (0.001 m N  $\approx$  14 ppm by mass N) nitrogen concentrations. Unreactive aqueous nitrogen ( $\text{N}_2^0 + \text{NH}_3^0$ ) progressively increase in N concentration and then dominate over reactive nitrogen ( $\text{NH}_4^+$ ) in fluids in equilibrium with the eclogite-facies assemblage while temperature increases from 600–1000 °C (Fig. 2a–b). For the model peridotite assemblage the relationship is similar for high nitrogen concentrations, where molecular aqueous nitrogen ( $\text{N}_2$ ) progressively increases in abundance and then dominates over reactive nitrogen ( $\text{NH}_4^+$ ), but impor-

tantly, ammonia is always less dominant than what we find for the eclogite-facies eclogites assemblage (Fig. 3b). However, for low N concentrations in equilibrium with the peridotite assemblage, ammonium is the most abundant species across all temperatures investigated (Fig. 3a). Therefore, the hottest subduction zone fluids at QFM-2 are predicted to be dominated by  $\text{N}_2$  at higher N concentrations in both eclogite-facies (Fig. 2b) and model peridotite (Fig. 3b) assemblages, but at the lowest N-concentrations,  $\text{N}_2$  is minor for both eclogite (Fig. 2a) and peridotite (Fig. 3a) with ammonia and ammonium dominating, respectively. Note, the difference in domination for ammonia and ammonium between the eclogite and peridotite simulations is the function of differing pH values for the fluids at high temperatures (Fig. 1).

## 4. DISCUSSION

### 4.1. Limitations of the approach

This study examines the role of pH in the deep-Earth nitrogen cycle at temperatures from 400 to 1000 °C and pressures of 1–5 GPa (Figs. 1–3). We have calculated the pH for aqueous fluids in equilibrium with hypothetical eclogite-facies and metasediments with a simple bulk chemistry. However, our approach does not consider some important and classically considered electron-donor elements (i.e.,  $\text{Fe}^{2+,3+}$ ,  $\text{S}^{-2,+6}$ ) nor do we consider the role of alkali metals besides  $\text{Na}^+$  (i.e.,  $\text{K}^+$ ,  $\text{Rb}^+$ ,  $\text{Cs}^+$ ). Furthermore, under water-saturated conditions at 1000 °C and 1 GPa, peridotites and eclogites would likely melt (Grove et al., 2006). The DEW model does not (presently) take melting into account, and therefore we can only show the predicted speciation of aqueous nitrogen in equilibrium with solid mineral phases. If a melt phase were present, the partitioning of ions could be affected, and therefore the pH values of the aqueous fluids could also be affected. Therefore, we acknowledge that our approach represents a simplification. Nonetheless, our approach does allow for robust constraints to be applied based on the effect of pH on the speciation of aqueous nitrogen at high temperatures and pressures. The modeling detailed here shows how changes in pH as a function of mineralogy are not only possible, but are predicted to be large (orders of magnitude – Fig. 1a–d), and this approach has been further reinforced by a recent study employing a more complex (theoretical) modeling approach (Galvez et al., 2016). Ergo, large changes in  $f\text{O}_2$  are predicted, which is consistent with empirical data and conventional understanding. However, we also predict large variations in pH, which also overlap with speciation changes for nitrogen between three distinct species ( $\text{NH}_4^+$ - $\text{NH}_3^0$ - $\text{N}_2^0$ ).

### 4.2. Ammonia and ammonium

Our results show that nitrogen can occur either in a neutral or positively charged state at high temperatures in the mantle, where molecular vs. ammoniac is dependent on  $f\text{O}_2$  (i.e. Mikhail and Sverjensky, 2014), but ammonia vs. ammonium is dependent on pH (this study). For example,




Fig. 2. Aqueous speciation of nitrogen in fluids in equilibrium with a model eclogite-facies mineral assemblages at 5 GPa, and QFM -2, and temperatures ranging from cold to hot subduction zone conditions. (a) Jadeite + Pyrope + Kyanite + diamond in equilibrium with fluid containing 0.001 m N, and (b) Jadeite + Pyrope + Kyanite + diamond in equilibrium with fluid containing 0.1 m N.

at 700 °C, 5 GPa, and a  $\log f_{\text{O}_2} - 2$  ( $\Delta\text{QFM}$ ) > 80% of the nitrogen in equilibrium with Jadeite + Pyrope + Kyanite + Coesite + Diamond will be  $\text{NH}_4^+$  but neutrally charged nitrogen becomes dominant with increasing temperature, where  $\text{NH}_4^+$  drops off to 10% and  $\text{N}_2^0$  and  $\text{NH}_3^0$  make up 60% and 30% of the fluid, respectively, at 1000 °C (Fig. 2a). At 1000 °C and 5 GPa (Fig. 1d), aqueous nitrogen in fluids under equilibrium with eclogite are predicted to be more alkaline than in peridotite, and there are a wide range of conditions where  $\text{NH}_3^0$  will be the dominant nitrogen species (Fig. 1d). As a further additional complexity influencing the behavior of nitrogen in upper mantle fluids, the total amount of nitrogen in the fluids also affects the

speciation (Figs. 2 and 3). For example, in Fig. 1a, the speciation of nitrogen in eclogitic fluids at QFM - 2, ranges from  $\text{NH}_3^0$  in low-N fluids to  $\text{N}_2$  in high-N fluids.

We note that the dominance of  $\text{NH}_3^0$  in our hypothetical eclogite-facies and metasediments must be considered a gross simplification for the sake of constraining the first-order level effect of pH. Nonetheless, this result can be viewed as an upper limit, where the addition of  $\text{H}^+$  would protonate  $\text{NH}_3^0$  to  $\text{NH}_4^+$ . We further note that there are other reactions (in nature) which can produce  $\text{H}^+$  ions and convert  $\text{NH}_3^0$  to  $\text{NH}_4^+$ . For example, such reactions require coupling with a reaction that produces  $\text{H}^+$  ions during subduction (e.g., the carbonation of calcium shown in Eq. (6)):





Fig. 3. Aqueous speciation of nitrogen in fluids in equilibrium with a model peridotite mineral assemblages at 5 GPa, QFM -2, and temperatures ranging from cold to hot subduction zone conditions. (a) Forsterite + Enstatite + Pyrope/Clinochlore + diamond in equilibrium with fluid containing 0.001 m N, and (b) Forsterite + Enstatite + Pyrope/Clinochlore + diamond in equilibrium with fluid containing 0.1 m N. Note, the minerals are pyrope at 800 °C and above, clinochlore at 600 and 700 °C (data from Mikhail and Sverjensky, 2014).



Therefore, we can model the incorporation of neutrally charged ammonia into K-bearing minerals through pH-dependent reactions such as:



Because some carbonation reactions generate  $\text{H}^+$  ions (e.g. Eq. (6)), the pH of fluids in equilibrium with eclogites may not inherently destabilize ammonium, but instead the stability of ammonium over ammonia in nature may depend on the bulk chemistry of the more complex natural system(s). Therefore, future empirical work may indeed link the carbon cycle with the nitrogen cycle, on a more genetic

basis. To take this further requires extensive experimental and theoretical interrogation in the near future to examine the pH shift and the dramatic effect on N speciation (shown here). However, where conditions permit and ammonium is stabilized, the secondary  $\text{H}^+$ -producing reactions (e.g., Eq. (6)) are not required for nitrogen to exchange for potassium directly (Eq. (8)):



Our data also show that fluid mobilization from the sedimentary package into the mantle (eclogite or peridotite) will result in transitions between  $\text{N}_2^0$ ,  $\text{NH}_3^0$ , and  $\text{NH}_4^+$ , where the nature of these transitions will vary depending on whether




Fig. 4. A cartoon showing the likely pathways followed by nitrogen during subduction. This cartoon illustrates that there are two options for nitrogen, degassing of neutrally charged  $\text{NH}_3^0$  and  $\text{N}_2^0$ , or re-gassing of the mantle with lattice-bound  $\text{NH}_4^+$  transported by the mineralogical conveyor belt of K-bearing minerals shown as colored arrows (phengite, phlogopite, K-bearing clinopyroxene, Phase-X and K-Hollandite). The depth-related stability for the K-bearing phases shown were taken from Harlow and Davies (2004). (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

or not the reaction is driven by pH,  $f\text{O}_2$ , or even the N concentration in the fluid (Figs. 2 and 3). In short, the fluid pathway and bulk compositions (i.e., slab-mantle systems, slab-ambient mantle, and mantle wedge-ambient mantle) will determine the aqueous N-speciation (Fig. 1a–d). Therefore, it is conceivable that nitrogen can behave like a highly incompatible and volatile element (e.g., a noble gas) or a large-ion lithophile element (e.g.,  $\text{K}^+$  &  $\text{Rb}^+$ ) in the same dynamic upper-mantle wedge system (see Fig. 4). This means some arc systems can out-gas + in-gas, some will solely in-gas, and others will solely out-gas nitrogen depending upon the bulk rock geochemistry of the system. We argue that these data strongly imply that cold subduction zones favor mass transfer of N into the mantle, but hot subduction zones do not (Fig. 2). This notion partly explains why efforts to determine a global N flux using specific geographical localities have been thwarted by contradictory (but equally correct) results (e.g., Fischer et al., 2002; Busigny et al., 2003; Elkins et al., 2006; Barry and Hilton, 2016). In fact, we argue that contrasting results should actually be expected.

In short, we propose that nitrogen should not have a single label regarding geochemical behavior. Nitrogen should not be considered lithophile, siderophile, or volatile, but instead nitrogen should always be viewed as a most dynamic, *chameleon* element whose behavior in aqueous fluids is effectively determined by a combination of temperature, pressure, oxygen fugacity, pH, and N concentration (i.e. mole fraction).

## 5. BROADER IMPLICATIONS

### 5.1. Nitrogen isotope fractionation

The results discussed above have implications for the isotopic evolution of subducted nitrogen during devolatilization of the slab and/or mantle, because the magnitude and direction of  $\Delta^{15}\text{N}$  for  $\text{NH}_3^0\text{-N}_2^0$ ,  $\text{NH}_4^+\text{-N}_2^0$ , and  $\text{NH}_4^0\text{-NH}_3^0$  differ dramatically. For example, at 600 °C the

predicted  $\Delta^{15}\text{N}_{\text{NH}_3\text{-N}_2}$  is  $-4\%$ ,  $\Delta^{15}\text{N}_{\text{NH}_4\text{-N}_2}$  is  $+2\%$ , and  $\Delta^{15}\text{N}_{\text{NH}_4\text{-NH}_3}$  is  $+8\%$  (Hanschmann, 1981). Therefore, the evolution of the  $\delta^{15}\text{N}$  value of subducted nitrogen cannot be modeled with a single fractionation factor ( $\Delta^{15}\text{N}$ ), because during progressive devolatilization of the slab in the mantle the magnitude and direction of  $\Delta^{15}\text{N}$  depends on upon the coupled  $f\text{O}_2$ -pH conditions for a given temperature, which should not be considered uniform across different mantle-wedge systems. Furthermore, because the stability of the K-bearing phases is a function of P-T- $X_{\text{K}}$ , it is unlikely to be constant with depth on a global scale (because the large P-T variability of arc systems globally; Syracuse et al., 2010). Therefore, the magnitude and direction of  $\Delta^{15}\text{N}$  during devolatilization of nitrogen from the slab or mantle is difficult to constrain. Importantly, if the reaction in question is  $\Delta^{15}\text{N}_{\text{NH}_3\text{-N}_2}$  where the direction is for a  $^{15}\text{N}$ -depleted residuum, this may explain why some eclogitic diamonds show mantle-like negative  $\delta^{15}\text{N}$  values alongside crustal organic carbon-like light  $\delta^{13}\text{C}$  values (Cartigny et al., 1997, 1998). In addition, these data could speak to why mantle diamonds show a much larger range of nitrogen isotope values relative to carbon (Mikhail et al., 2014) alongside disparate nitrogen abundance trends (Mikhail & Howell, 2016).

### 5.2. The pH of mantle fluids

In low-temperature geochemistry, Eh and pH are both considered important variables for predicting and expressing the nature of given chemical environments (i.e., for a recent review on low-T nitrogen see Stüeken et al., 2016). Traditionally, only Eh is considered in high temperature geochemistry, and because mineral charge-balances reflect the electron exchange where most minerals receive their negative charge in the form of  $\text{O}^-$  anions, this parameter is commonly expressed as the fugacity of oxygen ( $f\text{O}_2$ ). Furthermore, traditional models for the speciation of volatile elements in mantle fluids have long been constructed based on mixtures of neutral gases ( $\text{CO}_2$ ,  $\text{CH}_4$ ,  $\text{H}_2$  and  $\text{H}_2\text{O}$ ; com-

monly termed COH-fluids; e.g. Zhang and Duan, 2009). Thus, the role of dissolved aqueous ions or species derived from silicate rock components has been overlooked or ignored (see Sverjensky and Huang, 2015). Because oxygen fugacity is considered of primary importance there have been numerous theoretical and empirical approaches to predicting and quantifying the fugacity of oxygen in the mantle during accretion and differentiation (Wade and Wood, 2005; Wood et al., 2006; Frost et al., 2008), in upper mantle peridotites (Wood and Virgo, 1989; Canil et al., 1990; Woodland et al. 1992, 1996, 2006; Ionov and Wood 1992 Brandon and Draper 1996; Rohrbach et al., 2007; Cottrell and Kelley, 2013), lower mantle peridotites (Frost et al., 2004), in eclogites (Simakov, 2006; Stagno et al., 2015; Smart et al., in press), in the mantle wedge of arc systems (Wood et al., 1990; Lécuyer and Ricard, 1999; Parkinson and Arculus, 1999), and within other telluric bodies in the Solar System (Herd, 2008).

DEW model predictions now suggest pH should also be considered a significant dimension in high temperature mineralogical composition and mantle geochemistry (Mikhail and Sverjensky, 2014; Sverjensky et al., 2014b; Sverjensky and Huang, 2015; and this study), although other modeling approaches are being pioneered that have led to the same conceptual conclusion (Galvez et al., 2016). In hindsight, the importance of high temperature pH is not a surprising result. It is well established that at lower temperatures (<ca. 400 °C) fluid-rock interaction can exert large shifts in fluid pH, which can have dramatic effects. For example, pH as a variable has been explored by experimental means to understand why some economically viable REE deposits show significant REE fractionation leading to economically viable HREE-enrichment (Migdisov et al., 2009; Williams-Jones et al., 2012).

The most important finding of this contribution is the expected variability of the pH values of fluids in equilibrium with peridotite- and eclogite-facies assemblages across several units, even for temperatures up to 1000 °C (Fig. 1b and d). In the case of carbon (Sverjensky et al., 2014b; Galvez et al., 2016) and nitrogen (this study) the large pH shift as a function of mineral chemistry will significantly affect the speciation and behavior of these important elements (in the mantle and in subduction zones), and should therefore have influenced which species of pH sensitive compounds were and are degassed into planetary surface environments. Finally, our theoretical study raises several pressing questions:

- (1) Is there stratification of aqueous fluid pH with depth in the bulk silicate Earth following pressure effects on  $Mg^{2+}/Na^{+}$  phase equilibria and phase transitions (coesite- $\alpha$ -Qtz- $\beta$ -SiO<sub>2</sub>; Fig. 1a and b)?
- (2) How much (if at all) has the pH of aqueous mantle fluids changed through time?
- (3) How does the pH of aqueous mantle fluids vary between the different inner solar system planets (as is the case for  $fO_2$ )?
- (4) How accurate is the large pH shift as a function of mineralogical composition predicted in this study (Fig. 1a–d)?

- (5) Which reactions or elemental fractionations can be used to retrospectively constrain the pH of a metasomatized high temperature environment using data from silicate/oxide minerals or melt inclusions?

## 6. CONCLUSIONS

We have calculated the speciation of aqueous nitrogen in equilibrium with a model eclogite-facies mafic mineral assemblage and a model carbonaceous eclogite-facies mafic mineral assemblage using the DEW model. We find nitrogen to be an extremely dynamic element whose speciation and behavior is determined by a combination of temperature, pressure, oxygen fugacity, N concentration, and pH. Our modeling results show that increasing temperature stabilizes molecular nitrogen and increasing pressure stabilizes ammoniac nitrogen – but the nitrogen concentration also exerts a governing control (Figs. 2 and 3). These show that the pH of aqueous fluids is controlled by mineralogy for a given pressure and temperature and can vary by up to 4 units at 5 GPa and 1000 °C (Fig. 1). This finding clearly demonstrates that pH plays an important role in controlling speciation, and thus mass transport, of Eh-pH sensitive elements (such as nitrogen) up to at least 1000 °C.

## ACKNOWLEDGEMENTS

SM is grateful to the Carnegie Institution of Washington for funding this work through the bestowment of a Carnegie Postdoctoral Fellowship. SM also acknowledges the School of Earth and Environmental Science (St Andrews) for providing a start-up fund which assisted in the development of these data. DAS is grateful to grants from the Sloan Foundation through the Deep Carbon Observatory (Reservoirs and Fluxes, and Extreme Physics and Chemistry programs) and a community-building Officer Grant from the Sloan Foundation, as well as support from NSF grants EAR-1624325 and EAR-1624325, and a grant from the W.M. Keck Foundation (The Co-Evolution of the Geo- and Biosphere) Grant #10583-02 to Sverjensky. PHB's contribution was supported in part by the NSF grant EAR-1144559 (A Petrological and N Isotope Study Of Crustal Recycling Through Time). We wish to acknowledge discussions with David Hilton and Tobias Fischer which assisted the development of this contribution. We also wish to thank Associate Editor Prof Horst Marschall, Dr Ralf Halama (Keele University) and two anonymous reviewers for their thoughtful comments and criticisms which improved the overall quality of this manuscript. This study directly benefitted from the financial support of the Deep Carbon Observatory via attendance at the Second Early Career Scientist Workshop in the Azores (SM and PHB). Some of the ideas presented in this paper were developed and refined during their time together as room mates on the welcoming, small, and beautiful volcanic island of São Miguel. A noteworthy show of gratitude is also provided for the workshops organising committee: Donato (The Don) Giovannelli, Fátima Viveiros, Alyssia Cox, Cody Sheik, Daniel (Humvee) Hummer, Dana Thomas, Katie Pratt.

## REFERENCES

- Ague J. J. and Nicolescu S. (2014) Carbon dioxide released from subduction zones by fluid-mediated reactions. *Nat. Geosci.* **7**, 355–360.

- Andersen T., Burke E. A. J. and Neumann E. (1995) Nitrogen-rich fluid in the upper mantle: fluid inclusions in spinel dunite from Lanzarote, Canary Islands. *Contrib. Miner. Petrol.* **120**, 20–28.
- Barry P. H., Hilton D. R., Halldórsson S. A., Hahn D. and Marti K. (2012) High precision nitrogen isotope measurements in oceanic basalts using a static triple collection noble gas mass spectrometer. *Geochem. Geophys. Geosyst.* **13**(1).
- Barry P. H. and Hilton D. R. (2016) Release of subducted sedimentary nitrogen throughout Earth's mantle. *Geochem. Perspect. Lett.* **2**, 148–159.
- Bebout G. E., Fogel M. L. and Cartigny P. (2013) Nitrogen: highly volatile yet surprisingly compatible. *Elements* **9**, 333–338.
- Bebout G., Lazzeri K. and Geiger C. A. (2016) Pathways for nitrogen cycling in Earth's crust and upper mantle: a review and new results for microporous beryl and cordierite. *Am. Miner.* **101**, 7–24.
- Blundy J. and Wood B. J. (2003) Partitioning of trace elements between crystals and melts. *Earth Planet. Sci. Lett.* **210**, 383–397.
- Brandon A. D. and Draper D. S. (1996) Constraints on the origin of the oxidation state of mantle overlying subduction zones: an example from Simcoe, Washington, USA. *Geochim. Cosmochim. Acta* **60**, 1739–1749.
- Brooker R. A., Du Z., Blundy J., Kelley S. P., Allan N. L., Wood B. J., Chamorro E. M., Wartho J. A. and Purton J. A. (2003) The 'zero charge' partitioning behaviour of noble gases during mantle melting. *Nature* **423**, 738–741.
- Busigny V. and Bebout G. E. (2013) Nitrogen in the silicate Earth: speciation and isotopic behaviour during mineral-fluid interactions. *Elements* **9**, 353–358.
- Busigny V., Cartigny P., Philippot P., Ader M. and Javoy M. (2003) Massive recycling of nitrogen and other fluid-mobile elements (K, Rb, Cs, H) in a cold slab environment: evidence from HP to UHP oceanic metasediments of the Schistes Lustrés nappe (western Alps, Europe). *Earth Planet. Sci. Lett.* **215**, 27–42.
- Busigny V., Cartigny P. and Philippot P. (2011) Nitrogen isotopes in ophiolitic metagabbros: a re-evaluation of modern nitrogen uxes in subduction zones and implication for the early Earth atmosphere. *Geochim. Cosmochim. Acta* **75**, 7502–7521.
- Busigny V., Lebeau O., Ader M., Krapez B. and Bekker A. (2013) Nitrogen cycle in the late archaean ferruginous ocean. *Chem. Geol.* **362**, 115–130.
- Canil D., Virgo D. and Scarfe C. M. (1990) Oxidation state of mantle xenoliths from British Columbia, Canada. *Contrib. Miner. Petrol.* **104**, 453–462.
- Cartigny P., Boyd S. R., Harris J. W. and Javoy M. (1997) Nitrogen isotopes in peridotitic diamonds from Fuxian, China: the mantle signature. *Terra Nova* **9**, 175–179.
- Cartigny P., Harris J. W. and Javoy M. (1998) Eclogitic diamond formation at Jwaneng: no room for a recycled component. *Science* **280**, 1421–1424.
- Cottrell E. and Kelley K. A. (2013) Redox heterogeneity in mid-ocean ridge basalts as a function of mantle source. *Science* **340**, 1314–1317.
- Dauphas N. and Marty B. (1999) Heavy nitrogen in carbonatites of the Kola Peninsula: a possible signature of the deep mantle. *Science* **286**, 2488–2490.
- Dobrzhinetskaya L. F., Wirth R., Yang J., Hutcheon I. D., Weber P. K. and Green H. W. (2009) High-pressure highly reduced nitrides and oxides from chromitite of a Tibetan ophiolite. *Proc. Natl. Acad. Sci.* **106**, 19233–19238.
- Elkins L. J., Fischer T. P., Hilton D. R., Sharp Z. D., McKnight S. and Walker J. (2006) Tracing nitrogen in volcanic and geothermal volatiles from the Nicaraguan volcanic front. *Geochim. Cosmochim. Acta* **70**, 5215–5235.
- Facq S., Daniel I. and Sverjensky D. A. (2014) *In situ* Raman study and thermodynamic model of aqueous carbonate speciation in equilibrium with aragonite under subduction zone conditions. *Geochim. Cosmochim. Acta* **132**, 375–390.
- Fischer T. P., Hilton D. R., Zimmer M. M., Shaw A. M., Sharp Z. D. and Walker J. A. (2002) Subduction and recycling of nitrogen along the Central American margin. *Science* **297**, 1154–1157.
- Fischer T. P., Takahata N., Sano Y., Sumino H. and Hilton D. R. (2005) Nitrogen isotopes of the mantle: insights from mineral separates. *Geophys. Res. Lett.* **32**. <http://dx.doi.org/10.1029/2005GL022792>.
- Frost D. J. and McCammon C. M. (2008) The redox state of earth's mantle. *Annual Rev. Earth Planet. Sci.* **36**, 389–420.
- Frost D. J., Liebske C., Langenhorst F., McCammon C. A. and Trønnes R. G. (2004) Experimental evidence for the existence of iron-rich metal in the Earth's lower mantle. *Nature* **428**, 409–412.
- Frost D. J., Mann U., Asahara Y. and Rubie D. C. (2008) The redox state of the mantle during and just after core formation. *Phil. Trans. R. Soc. A* **366**, 4315–4337.
- Galvez M. E., Connolly J. A. D. and Manning C. E. (2016) Implications for metal and volatile cycles from the pH of subduction zone fluids. *Nature* **539**, 420–424.
- Grove T. L., Chatterjee N., Parman S. W. and Medard E. (2006) The influence of H<sub>2</sub>O on mantle wedge melting. *Earth Planet. Sci. Lett.* **249**, 74–89.
- Haendel D., Muhle K., Nitzsche H. M., Stiehl G. and Wand U. (1986) Isotopic variations of the fixed nitrogen in metamorphic rocks. *Geochim. Cosmochim. Acta* **50**, 749–758.
- Halama R., Bebout G., John T. and Scambelluri M. (2014) Nitrogen recycling in subducted mantle rocks and implications for the global nitrogen cycle. *Int. J. Earth Sci.* **103**, 2081–2099.
- Halama R., Bebout G. E., John T. and Schenk V. (2010) Nitrogen recycling in subducted oceanic lithosphere: the record in high- and ultrahigh-pressure metabasaltic rocks. *Geochim. Cosmochim. Acta* **74**, 1636–1652.
- Hanschmann G. (1981) Berechnung von Isotopieeffekten auf quantenchemischer Grundlage am Beispiel stickstoffhaltiger Moleküle. *ZFI Mitteilungen* **41**, 19–31.
- Harlow G. E. and Davies R. (2004) Status report on stability of K-rich phases at mantle conditions. *Lithos* **77**, 647–653.
- Helgeson H. C. and Kirkham D. H. (1974a) Theoretical prediction of the thermodynamic behavior of aqueous electrolytes at high pressures and temperatures: I. Summary of the thermodynamic/electrostatic properties of the solvent. *Am. J. Sci.* **274**, 1089–1198.
- Helgeson H. C. and Kirkham D. H. (1974b) Theoretical prediction of the thermodynamic behavior of aqueous electrolytes at high pressures and temperatures: II. Debye-Hückel parameters for activity coefficients and relative partial molal properties. *Am. J. Sci.* **274**, 1199–1261.
- Helgeson H. C. and Kirkham D. H. (1976) Theoretical prediction of the thermodynamic properties of aqueous electrolytes at high pressures and temperatures. III. Equation of state for aqueous species at infinite dilution. *Am. J. Sci.* **276**, 97–240.
- Helgeson H. C., Kirkham D. H. and Flowers G. C. (1981) Theoretical prediction of the thermodynamic behavior of aqueous electrolytes at high pressures and temperatures. IV. Calculation of activity coefficients, osmotic coefficients, and apparent molal and standard and relative partial molal properties to 5 kb and 600 C. *Am. J. Sci.* **281**, 1241–1516.
- Herd C. D. K. (2008) Basalts as probes of planetary interior redox state. In *Oxygen in the Solar System* (eds. G. J. MacPherson, D. W. Mittlefehldt and J. H. Jones). Mineralogical Society of America, Chantilly, VA.

- Honma H. and Itihara Y. (1983) Distribution of ammonium in minerals of metamorphic and granitic rocks. *Geochim. Cosmochim. Acta* **45**, 983–988.
- Ionov D. A. and Wood B. J. (1992) The oxidation state of subcontinental mantle: oxygen thermobarometry of mantle xenoliths from central Asia. *Contrib. Mineral. Petrol.* **111**, 179–193.
- Johnson B. and Goldblatt C. (2015) The nitrogen budget of Earth. *Earth Sci. Rev.* **148**, 150–173.
- Johnson J. W., Oelkers E. H. and Helgeson H. C. (1992) SUPCRT92: a software package for calculating the standard molal thermodynamic properties of minerals, gases, aqueous species, and reactions from 1 to 5000 bars and 0 to 1000 °C. *Comput. Geosci.* **18**, 899–947.
- Lécuyer C. and Ricard Y. (1999) Long-term fluxes and budget of ferric iron: implication for the redox states of the Earth's mantle and atmosphere. *Earth Planet. Sci. Lett.* **165**, 197–211.
- Li Y. and Keppler H. (2014) Nitrogen speciation in mantle and crustal fluids. *Geochim. Cosmochim. Acta* **129**, 13–32.
- Li Y., Wiedenbeck M., Shcheka S. and Keppler H. (2013) Nitrogen solubility in upper mantle minerals. *Earth Planetary Sci. Lett.* **377**, 311–323.
- Marty B. and Dauphas N. (2003) The nitrogen record of crust-mantle interaction and mantle convection from Archean to present. *Earth Planet. Sci. Lett.* **206**, 397–410.
- Marty B., Zimmermann L., Pujol M., Burgess R. and Philippot P. (2013) Nitrogen isotopic composition and density of the Archean atmosphere. *Science* **342**, 101–104.
- Marty B. (1995) Nitrogen content of the mantle inferred from N<sub>2</sub>-Ar correlation in oceanic basalts. *Nature* **377**, 326–329.
- Migdisov A. A., Williams-Jones A. E. and Wagner T. (2009) An experimental study of the solubility and speciation of the Rare Earth Elements (III) in fluoride- and chloride-bearing aqueous solutions at temperatures up to 300 °C. *Geochim. Cosmochim. Acta* **73**, 7087–7109.
- Mikhail S. and Howell D. (2016) A petrological assessment of diamond as a recorder of the mantle nitrogen cycle. *Am. Mineral.* **101**, 780–787.
- Mikhail S. and Sverjensky D. A. (2014) Nitrogen speciation in upper mantle fluids and the origin of Earth's nitrogen-rich atmosphere. *Nat. Geosci.* **7**, 816–819.
- Mikhail S., Verchovsky A. B., Howell D., Hutchison M. T., Southworth R., Thomson A. R., Warburton P., Jones A. P. and Milledge H. J. (2014) Constraining the internal variability of the stable isotopes of carbon and nitrogen within mantle diamonds. *Chem. Geol.* **366**, 14–23.
- Mohapatra R., Harrison D., Ott U., Gilmour J. and Trierloff M. (2009) Noble gas and nitrogen isotopic components in Oceanic Island Basalts. *Chem. Geol.* **266**, 29–37.
- Palot M., Cartigny P., Harris J. W., Kaminsky F. V. and Stachel T. (2012) Evidence for deep mantle convection and primordial heterogeneity from nitrogen and carbon stable isotopes in diamond. *Earth Planet. Sci. Lett.* **357–358**, 179–193.
- Palya A. P., Buick I. S. and Bebout G. E. (2011) Storage and mobility of nitrogen in the continental crust: evidence from partially melted metasedimentary rocks, Mt. Stafford, Australia. *Chem. Geol.* **281**, 211–226.
- Pan D., Spanu L., Harrison B., Sverjensky D. A. and Galli G. (2013) The dielectric constant of water under extreme conditions and transport of carbonates in the deep Earth. *Proc. Nat. Acad. Sci.* **110**, 6646–6650.
- Parkinson I. J. and Arculus R. J. (1999) The redox state of subduction zones: insights from arc-peridotites. *Chem. Geol.* **160**, 409–423.
- Philippot P., Busigny V., Scambelluri M. and Cartigny P. (2007) Oxygen and nitrogen isotopes as tracers of fluid activities in serpentinites and metasediments during subduction. *Mineral. Petrol.* **91**, 11–24. <http://dx.doi.org/10.1007/s00710-007-0183-7>.
- Rohrbach A., Ballhaus C., Golla-Schindler U., Ulmer P., Kamenetsky V. S. and Kuzmin D. V. (2007) Metal saturation in the upper mantle. *Nature* **449**, 456–458.
- Shock E. L., Oelkers E. H., Johnson J. W., Sverjensky D. A. and Helgeson H. C. (1992) Calculation of the thermodynamic and transport properties of aqueous species at high pressures and temperatures: effective electrostatic radii to 1000 °C and 5 kb. *Faraday Soc. Trans.* **88**, 803–826.
- Shock E. L. and Helgeson H. C. (1988) Calculation of the thermodynamic and transport properties of aqueous species at high pressures and temperatures: correlation algorithms for ionic aqueous species and equation of state predictions to 5 kb and 1000 °C. *Geochim. Cosmochim. Acta* **52**, 2009–2036.
- Simakov S. R. (2006) Redox state and peridotites from sub-cratonic upper mantle and a connection with diamond genesis. *Contrib. Mineral. Petrol.* **151**, 282–296.
- Smart K. S., Chacko T., Stachel T., Muehlenbachs K., Stern R. A. and Heaman L. M. (2011) Diamond growth from oxidized carbon sources beneath the Northern Slave Craton, Canada: a  $\delta^{13}\text{C}$ -N study of eclogite-hosted diamonds from the Jericho kimberlite. *Geochim. Cosmochim. Acta* **75**, 6027–6047.
- Smart, K.S., Tappe, S., Simonetti, A., Simonetti, S.S., Woodland, A.B., Harris, C., in press. Tectonic significance and redox state of Paleoproterozoic eclogite and pyroxenite components in the Slave cratonic mantle lithosphere, Voyageur kimberlite, Arctic Canada. *Chem. Geol.* doi: <http://dx.doi.org/10.1016/j.chemgeo.2016.10.014>.
- Smith E. M., Kopylova M. G., Frezzotti M. L. and Afanasiev V. P. (2014) N-rich fluid inclusions in octahedrally-grown diamond. *Earth Planet. Sci. Lett.* **393**, 39–48.
- Som S. M., Buick R., Hagadorn J. W., Blake T. S., Perreault J. M., Harnmeijer J. P. and Catling D. C. (2016) Earth's air pressure 27 billion years ago constrained to less than half of modern levels. *Nature Geosci.* (advance online publication).
- Som S. M., Catling D. C., Harnmeijer J. P., Polivka P. M. and Buick R. (2012) Air density 2.7 billion years ago limited to less than twice modern levels by fossil raindrop imprints. *Nature* **484**, 359–362.
- Stagno V., Frost D. J., McCammon C. A., Mohseni H. and Fei Y. (2015) The oxygen fugacity at which graphite or diamond forms from carbonate-bearing melts in eclogitic rocks. *Contrib. Mineral. Petrol.* **169**, 16.
- Stüeken E. A., Kipp M. A., Koehler M. C. and Buick R. (2016) The evolution of Earth's biogeochemical nitrogen cycle. *Earth Sci. Rev.* **160**, 220–239.
- Sverjensky D. A. and Huang F. (2015) Diamond formation due to a pH drop during fluid-rock interactions. *Nature Commun.* **6**, 8702.
- Sverjensky D. A., Harrison B. and Azzolini D. (2014a) Water in the deep Earth: The dielectric constant and the solubilities of quartz and corundum to 60 kb and 1200 °C. *Geochim. Cosmochim. Acta* **129**, 125–145.
- Sverjensky D. A., Stagno V. and Huang F. (2014b) Important role for organic carbon in subduction-zone fluids in the deep carbon cycle. *Nat. Geosci.* **7**, 909–913.
- Syracuse E. M., van Keken P. E. and Abers G. A. (2010) The global range of subduction zone thermal models. *Phys. Earth Planet. Inter.* **51**, 1761–1782.
- Tanger J. C. and Helgeson H. C. (1988) Calculation of the thermodynamic and transport properties of aqueous species at high pressures and temperatures: revised equations of state for the standard partial molal properties of ions and electrolytes. *Am. J. Sci.* **288**, 19–98.

- Wade J. and Wood B. J. (2005) Core formation and the oxidation state of the Earth. *Earth Planet. Sci. Lett.* **236**, 78–95.
- Watenphul A., Wunder B., Wirth R. and Heinrich W. (2010) Ammonium-bearing clinopyroxene: a potential nitrogen reservoir in the Earth's mantle. *Chem. Geol.* **270**, 240–248.
- Williams-Jones A. E., Migdisov A. A. and Samson I. M. (2012) Hydrothermal mobilization of the rare earth elements – a tale of “Cerium” and “Yttrium”. *Elements* **8**, 355–360.
- Wood B. J. and Virgo D. (1989) Upper mantle oxidation state: ferric iron contents of lherzolite spinels by  $^{57}\text{Fe}$  Mössbauer spectroscopy and resultant oxygen fugacities. *Geochim. Cosmochim. Acta* **53**, 1277–1291.
- Wood B. J., Bryndzia L. T. and Johnson K. E. (1990) Mantle oxidation state and its relationship to tectonic environment and fluid speciation. *Science* **248**, 337–345.
- Wood B. J., Walter M. J. and Wade J. (2006) Accretion of the earth and segregation of its core. *Nature* **441**, 825–833.
- Woodland A. B., Kornprobst J. and Wood B. J. (1992) Oxygen thermobarometry of orogenic lherzolite massifs. *J. Petrol.* **33**, 203–230.
- Woodland A. B., Kornprobst J., McPherson E., Bodinier J.-L. and Menzies M. A. (1996) Metasomatic interactions in the lithospheric mantle. Petrologic evidence from the Lherz massif, French Pyrenees. *Chem. Geol.* **134**, 83–112.
- Woodland A. B., Kornprobst J. and Tabit A. (2006) Ferric iron in orogenic lherzolite massifs and controls of oxygen fugacity in the upper mantle. *Lithos* **89**, 222–241.
- Yokochi R., Marty B., Chazot G. and Burnard P. (2009) Nitrogen in peridotite xenoliths: lithophile behaviour and magmatic isotope fractionation. *Geochim. Cosmochim. Acta* **73**, 4843–4861.
- Zerkle A. L. and Mikhail S. (2017) The geobiological nitrogen cycle: from microbes to the mantle. *Geobiology*, 1–10.
- Zhang C. and Duan Z. A. (2009) A model for C–O–H fluid in the Earth's mantle. *Geochim. Cosmochim. Acta* **73**, 2089–2102.

*Associate editor:* Horst R. Marschal